

Dark energy from gravitational collapse?

László Á. Gergely

*Departments of Theoretical and Experimental Physics, University of Szeged, Hungary**

(Dated: February 7, 2008)

Abstract

We discuss the status of both cosmological and black hole type singularities in the framework of the brane-world model of gravity. We point out that the Big Bang is not properly understood yet. We also show new features of the gravitational collapse on the brane, the most important being the production of dark energy during the collapse.

arXiv:gr-qc/0606073v1 16 Jun 2006

*Electronic address: gergely@physx.u-szeged.hu

Singularities, both cosmological and of black hole type, are intriguing objects predicted by general relativity. One of the most desired aims of quantum gravity is to avoid such infinities. Although interesting results stem out both from loop quantum gravity and string theory, no conclusive decision has been reached. In the absence of a final theory of quantum gravity the question arises how the recently introduced, modified, but still classical theories of gravity, motivated by string / M-theory, will tackle these topics.

One of the most promising such theory is the generalized Randall-Sundrum second model, introduced in its original form in [1], in which our observable universe is a 4-dimensional hypersurface (the brane), embedded into a 5-dimensional curved space-time (the bulk). Standard model fields live on the brane, which is held together by a tension λ . A positive brane tension allows to recover the Einstein-gravity on the brane, at low energies and insignificant bulk curvature.

When this scenario is employed as a cosmological model, the brane becomes a 4-dimensional Friedmann-Lemaître-Robertson-Walker (FLRW) universe. Cosmological evolution is consequence of the brane moving in the Schwarzschild - anti de Sitter bulk. The scale factor, related in a simple way to the distance from the bulk black hole, evolves driven by a *modified* Friedmann and a *modified* Raychaudhuri equation [2]. The modifications include a quadratic source term in the brane energy momentum (modifying early cosmology), a dark radiation term scaled by the mass of the bulk black hole (representing the electric part of the bulk Weyl curvature), source terms from a possible asymmetric embedding (which can cause late time acceleration) and the pull-back to the brane of any existing bulk energy momentum.

The Big Bang, occurring when the scale factor is zero, corresponds to the brane emerging from the singularity. However it would contradict basic principles of physics to allow the brane escaping from below the horizon. In consequence it is not clear, whether the brane somehow condensates outside the event horizon from (quantum) bulk fields, or the bulk black hole is created far away from the pre-existing brane. The latter possibility was investigated in [3], assuming a brane which is radiating into the fifth dimension. If so, the static character of the bulk is obstructed and its simplest model would be the Vaidya-anti de Sitter space-time. Cosmological evolution in this radiation-filled bulk was further exploited in [2]-[10]. The parameters of the model can be fitted in such a way that the late time universe is extremely close to what is predicted by general relativity and verified by astronomical observations.

Black holes are fascinating objects, first predicted by general relativity. The energy momentum, as source of gravity, curves space-time in the vicinity of these objects to such an extent, that even light rays can follow circular orbits, like the planets revolve about stars. Strangely enough, much closer to the center, the role of time and space is interchanged in some subtle way. The curvature of space-time approaches infinity near the center, forming a space-time singularity. According to the Cosmic Censorship hypothesis, an event horizon luckily hides the singularity from any outside viewer in the majority of cases. Even so, black holes are dangerous creatures. They accrete surrounding matter in galactic centers and force the magnetic field lines into configurations, which give birth to powerful jets. These are starring on some of the most astonishing astronomical pictures.

It is not surprising, that the search for black holes in the brane-world scenario has consumed much energy. But we face a series of negative results. In brane cosmological models there are black holes in the bulk, but not on the brane, where they are allowed just as minuscule test particles. Looking for spherically symmetric vacuum solutions on the brane, the Schwarzschild black hole turns out as the unique solution of the modified gravitational equations, provided (a) the bulk is empty (b) the Weyl curvature of the bulk has no electric part, and (c) the embedding of the brane is symmetric. However the singularity has to extend into the fifth dimension [11]. The emerging black string could decay into a black cigar, due to the Gregory-Laflamme instability [12]. However, more recently it was shown that under very mild assumptions, classical event horizons cannot pinch off [13].

Becoming a black string, the black hole singularity, resulted from the gravitational collapse of a stellar object on the brane leaves its cradle. Such a model, with black strings puncturing the cosmological brane was investigated in [14]. The brane shows a Swiss-cheese type structure, with Schwarzschild black "holes" in a "cheese" represented by the cosmological perfect fluid. In certain cases such a brane evolves towards a new type of *pressure singularity*, accompanied by a *regular* behavior of the cosmic acceleration.

By turning on the electric part of the bulk Weyl curvature, the modified Einstein equations are solved by the Reissner-Nordström black hole with mass m and tidal charge q [15]. The tidal charge arises from the Kaluza-Klein modes of gravity, as opposed to general relativity, where $q = Q^2$ represents the square of the electric charge Q . When $q > 0$, the tidal black hole is analogous to the charged black hole: it has two horizons, both below the Schwarzschild radius. A negative tidal charge, by contrast, increases the radius of the event

horizon. Gravity is strengthened on the brane, light deflection and gravitational lensing are stronger, than for a Schwarzschild black hole with the same mass. The bulk containing such a tidal black hole brane remains unknown.

Gravitational collapse of a dust sphere in general relativity results in a black hole. Does this phenomenon, known as the Oppenheimer-Snyder collapse [16] generalize to brane-worlds? The question was answered in [17]. The collapse of a homogeneous Kaluza-Klein energy density with static exterior leads to either a bounce, black hole formation or to a naked singularity. In the absence of the tidal charge the same analysis shows that the vacuum surrounding the collapsing sphere cannot be static.

This no-go result is based on the assumption that the collapsing matter is dust, but this can obviously be relaxed. We do this for the case of vanishing Kaluza-Klein modes. In other words, we consider a collapsing FLRW sphere with constant comoving radius χ_0 (and physical radius $R = \chi_0 a$), embedded into a Schwarzschild brane [19]. Switching out the tidal charge has better motivation as the desire for simplicity. Whenever the radius of the collapsing sphere is much higher than the characteristic scale of the extra dimension, the black-hole metric on the brane is Schwarzschild [20]. Whereas table-top experiments [21] on possible deviations from Newton's law currently probe gravity at submillimeter scales, this seems no severe restriction.

With a collapsing star composed of perfect fluid with energy density ρ and pressure p , the vacuum exterior with Schwarzschild mass m can be held static. For vanishing brane cosmological constant and flat spatial sections the generic junction conditions derived in [18] imply the following evolution of the scale factor a in comoving time τ :

$$a^{3/2} = a_0^{3/2} - \left(\frac{9m}{2\chi_0^3} \right)^{1/2} \tau . \quad (1)$$

Here a_0 represents the initial value of the scale factor. Inevitably, after comoving time $\tau_1 = (2\chi_0^3 a_0^3 / 9m)^{1/2}$, the collapse ends in the singularity $a = 0$.

The Schwarzschild mass is related to the "physical" mass $M = (4\pi R^3/3)\rho$ as

$$m = M \left(1 + \frac{\rho}{2\lambda} \right) . \quad (2)$$

How can m be interpreted from the interior, if different from the "physical" mass? As shown in [19], m represents the Bardeen quasilocal mass [22], summing up both matter and gravitational energy contributions. Both masses cannot be constant, as they differ by

a typical brane-world contribution ρ/λ (due to the source term quadratic in the energy momentum).

The Schwarzschild mass being held constant, M varies in a precise way during the collapse. The ensemble of the junction conditions and modified Einstein equations show that both ρ and p increase in magnitude towards infinite values [19]. Eq. (2) then implies that M should accordingly decrease, until it reaches zero. Where has the "physical" mass disappeared? We obtain the answer from the equation of state of the fluid:

$$\frac{p}{\lambda} = \frac{1}{2} \left(1 - \frac{\rho}{\lambda}\right) - \frac{1}{2} \left(1 + \frac{\rho}{\lambda}\right)^{-1}. \quad (3)$$

At the beginning of the collapse (when $\rho \ll \lambda$ holds) this approaches

$$p \approx -\frac{\rho^2}{2\lambda}. \quad (4)$$

This represents ordinary matter, satisfying all energy conditions. It is basically dust (a tiny negative pressure appears in the first order). Towards the final stages of collapse, when $\rho \gg \lambda$ Eq. (3) gives

$$p \approx -\frac{\rho}{2}. \quad (5)$$

The magnitude of the negative pressure has considerably increased such that the condition for dark energy $\rho + 3p \approx -\rho/2 < 0$ is satisfied. Thus *in the brane-world scenario the original matter configuration is turned into dark energy by gravitational collapse.*

The high negative pressure (tension) has the same effect as a positive cosmological constant. Otherwise stated, a tension is raising in the brane, slowing down gravitational contraction. In vain however, the contraction goes on until the singularity is formed. This is due to the quadratic source term, which dominates at high energy densities.

Note that in the general relativistic limit the tension vanishes, the fluid turns into dust and we recover the Oppenheimer-Snyder collapse.

We conclude with a few remarks. First, the notorious difficulty in finding black hole solutions on the brane may be related to the tensions arising in the gravitational collapse. Second, when black holes are formed on the brane, the initial configuration of ordinary matter surprisingly turns into dark energy. However, in brane-worlds even dark energy speeds up gravitational collapse. Third, a similar, but higher dimensional mechanism to what we have just described here may be responsible for the initial condensation of branes

(containing dark energy) from a primordial bulk. This would solve the Big Bang problem.

- [1] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 4690 (1999).
- [2] L.Á. Gergely, *Phys. Rev. D* **68**, 124011 (2003).
- [3] A. Chamblin, A. Karch, and A. Nayeri, *Phys. Lett. B*, **509**, 163 (2001).
- [4] D. Langlois, L. Sorbo, and M. Rodríguez-Martínez, *Phys. Rev. Lett.* **89**, 171301 (2002).
- [5] L.Á. Gergely, E. Leeper, and R. Maartens, *Phys. Rev. D* **70**, 104025 (2004).
- [6] D. Jennings, I. R. Vernon, *J. Cosmol. Astropartic. Phys.* JCAP **0507** 011 (2005).
- [7] D. Jennings, I.R. Vernon, A.-C. Davis, and C. van de Bruck, *J. Cosmol. Astropartic. Phys.* JCAP **0504** 013 (2005).
- [8] D. Langlois, *Is our Universe brany?*, hep-th/0509231 (2005).
- [9] L.Á. Gergely, Z. Keresztes, *J. Cosmol. Astropartic. Phys.* JCAP **06** (01), 022 (2006).
- [10] Z. Keresztes, I. Képiró, and L.Á. Gergely, *J. Cosmol. Astropartic. Phys.* JCAP **06** (05), 020 (2006).
- [11] A. Chamblin, S.W. Hawking, and H.S. Reall, *Phys. Rev. D* **61**, 065007 (2000).
- [12] R. Gregory, R. Laflamme, *Phys. Rev. Lett.* **70**, 2837 (1993).
- [13] G.T. Horowitz and K. Maeda, *Phys. Rev. Lett.* **87**, 131301 (2001).
- [14] L.Á. Gergely, *Brane-world cosmology with black strings*, to appear in *Phys. Rev. D*, hep-th/0603244 (2006).
- [15] N. Dadhich, R. Maartens, P. Papadopoulos, and V. Rezanian, *Phys. Lett B* **487**, 1 (2000).
- [16] J.R. Oppenheimer, H. Snyder, *Physical Review* **56**, 455 (1939).
- [17] M. Bruni, C. Germani, and R. Maartens, *Phys. Rev. Lett.* **87**, 231302 (2001).
- [18] L.Á. Gergely, *Phys. Rev. D* **71**, 084017 (2005), errata, *ibid.* **72**, 069902 (2005).
- [19] L.Á. Gergely, *Black holes and dark energy from gravitational collapse on the brane*, hep-th/0603254 (2006).
- [20] R. Emparan, G. T. Horowitz, and R. C. Myers, *Phys. Rev. Lett.* **85**, 499 (2000).
- [21] J.C. Long et al, *Nature* **421**, 922 (2003).
- [22] J.M. Bardeen, *Phys. Rev. Lett.* **46**, 382 (1981).